arXiv:1209.1054v1 [hep-ph] 5 Sep 2012

DCP-12-05

Symmetric texture-zero mass matrices and its eigenvalues

A. Criollo*
Área Académica de Matemáticas y Física,
Universidad Autónoma del Estado de Hidalgo,
Carr. Pachuca-Tulancingo Km. 4.5, C.P. 42184, Pachuca, Hgo.

R. Noriega-Papaqui†
Área Académica de Matemáticas y Física,
Universidad Autónoma del Estado de Hidalgo,
Carr. Pachuca-Tulancingo Km. 4.5, C.P. 42184, Pachuca, Hgo. and
Dual C-P Institute of High Energy Physics

Within the texture-zeros mechanism, first we provide necessary and sufficient conditions on the characteristic polynomial coefficients so that it has real, simple and positive roots, we traduce these conditions in terms to the invariants of the congruent matrices. Next all symmetric texture-zero mass matrices are counted and classified. Finally we apply in a systematic way the result from the first part to analyze the six, four and two zeros texture matrices presented in the second part.

PACS numbers: 12.5.Ff; 02.10.Ud

I. INTRODUCTION

In the Standard Model (SM) with $SU(2) \times U(1)$ as the gauge group of electroweak interactions, the mass physics of quarks and charged leptons are in the Yukawa Sector, after Spontaneous Symmetry Breaking (SSB), it is defined the mass matrix as:

$$M_f = \frac{v}{\sqrt{2}} Y_f, \qquad (f = u, d, l),$$

where v is the vacuum expectation value of the Higgs field and Y_f are the 3×3 Yukawa matrices. The physical masses of the particles are defined as the eigenvalues of the mass

---

*Electronic address: arturoc@uaeh.edu.mx

†Electronic address: rnoriega@uaeh.edu.mx

matrix M_f. Within the SM context the mass matrix is unknown, the only trail of the quarks mass matrices is the V_{ckm} matrix, that it is built by the product of matrices that diagonalize the u-quark mass matrix and d-quark mass matrix.

In 1977 Harald Fritzsch proposed a phenomenological study [1], the texture-zeros mechanism [1], that consist in looking for the simplest pattern of mass matrices, which can result self-consistent and experimentally reproduce the V_{ckm} parameters (elements).

Going in this direction, we present what kind of texture-zeros are well structured considering that by definition, the eigenvalues of the mass matrix are the masses of fermions or charged leptons, *i.e.*, they must be real, simple and positive. Mathematically speaking a symmetric mass matrix always guarantees that the physical masses are real, however, the positivity condition for the eigenvalues is not fulfilled by any symmetric matrix, moreover a positive definite matrix has eigenvalues real and positive, but no necessarily different.

The main result of this paper is to provide the rigorous mathematical structure for texture-zero matrices. In Sec. II, we show analytically how the mass matrices appears in the SM context. In Sec. III, we find necessary and sufficient conditions on the characteristic polynomial coefficients so that its roots are real, simple and positive. These conditions are rewriting in terms of the invariant congruent matrices, *i.e.* trace, determinant and trace of the power matrix. In Sec. IV, we develop a easy notation that counts and classifies the texture-zero matrices, and we show all symmetric matrices of 3×3 grouped into 2-zero, 4-zero and 6-zero texture, in order to complete the counting the matrix without zeros is included. Finally in Sec. V, we apply systematically the results of the Sec. III to all matrices of the Sec. IV. and we show what kind of texture matrices have real, different and positive eigenvalues and we present some numerical examples of these. For these examples the values of the masses are taken from web page PDG [2].

II. PRELIMINARIES

In the Yukawa sector of the SM, the mass terms for quarks and charged leptons can be expressed as

$$\bar{u}_L M_u u_R + \bar{d}_L M_d d_R + \bar{l}_L M_l l_R, \tag{1}$$

---

[1] For an excellent review see [3], and references there in.

where $u_{L(R)}$, $d_{L(R)}$ and $l_{L(R)}$ are the left(right)-handed quark and charged leptons fields for the u-sector (u, c, t), d-sector (d, s, b) and charged leptons (e, μ, τ) respectively. M_u, M_d and M_l are the mass matrices. Expressing the above equation in terms of the physical fields, one diagonalize the mass matrices by bi-unitary transformations

$$\begin{aligned} \bar{M}_u &= U^\dagger_{uL}\, M_u\, U_{uR} = Diag\,[m_u, m_c, m_t]\,, \\ \bar{M}_d &= U^\dagger_{dL}\, M_d\, U_{dR} = Diag\,[m_d, m_s, m_b]\,, \\ \bar{M}_l &= U^\dagger_{lL}\, M_l\, U_{lR} = Diag\,[m_e, m_\mu, m_\tau]\,, \end{aligned} \tag{2}$$

where U_{f_L} and U_{f_R} $(f = u, d, l)$ are in general complex unitary matrices. The quantities $m_u, m_d, \ldots$ etc. denote the eigenvalues of the mass matrices, i.e. the physical quark masses and they must have real and nonnegative quantities.

Re-expressing Eq. (1) in terms of physical fermion fields $(f'_{L(R)})$ as

$$\bar{u'}_L \bar{M}_u u'_R + \bar{d'}_L \bar{M}_d d'_R + \bar{l'}_L \bar{M}_l l'_R, \tag{3}$$

where $\bar{f'}_L = \bar{f}_L\, U_{fL}$ and $f'_R = U^\dagger_{fR}\, f_R$, $(f' = u', d', l')$.

Eq.(2) implies that $\bar{M}_f$ and M_f, $(f = u, d)$ are congruent matrices, the relation of congruence is an equivalence relation, which implies a space partition in cosets. Any two elements that belong at the same coset have the following invariants: determinant, trace, trace of the power matrix, characteristic polynomial and their eigenvalues, in the other hand, if $\bar{M}_f$ and M_f are congruent matrices then: $\det \bar{M}_f = \det M_f$, $\mathrm{tr}\, \bar{M}_f = \mathrm{tr}\, M_f$, $\mathrm{tr}\, \bar{M}_f^n = \mathrm{tr}\, M_f^n$, where n is a positive integer, $\det(\bar{M}_f - \lambda I) = \det(M_f - \lambda I)$ [2][4].

Considering to M_f as a 3×3 symmetric matrix with real coefficients then $\bar{M}_f$ is built as diagonal matrix where its elements are the eigenvalues of M_f, these eigenvalues are found as the roots its characteristic polynomial. The aim is find the necessary and sufficient conditions over the coefficients of the characteristic polynomial from M_f, *i.e* over the M_f elements, such that this polynomial has three real, positive and simple roots. This subject will be developed the following section.

---

[2] In this work, we will denote the product $(\mathrm{tr}\, A)(\mathrm{tr}\, A)$ as $\mathrm{tr}^2 A$. In the general case $(\mathrm{tr}\, A)^n = \mathrm{tr}^n A$ for n positive integer.

III. TEXTURES POLYNOMIAL

Definition 1 *A monic polynomial with real coefficients* $p(\lambda) = \lambda^3 + a_2\lambda^2 + a_1\lambda + a_0$, *is a texture polynomial if the roots of equation* $p(\lambda) = 0$ *are real, simple and positive, in other words, they are located in the positive semi-axis* $\mathbb{R}^+$ *and has multiplicity one.*

Lemma 1 *The polynomial of second degree* $p(\lambda) = \lambda^2 + a_1\lambda + a_0$ *has two real, simple and positive roots if and only if the following condition hold.*

$$a_1 < 0 < a_0 < \frac{a_1^2}{4}. \tag{4}$$

Proof. If $p(\lambda)$ has two real, simple and positive roots then there exists $\lambda_1, \lambda_2 \in \mathbb{R}^+,\ \lambda_1 \neq \lambda_2$ such that the polynomial can be expressed by

$$p(\lambda) = (\lambda - \lambda_1)(\lambda - \lambda_2) = \lambda^2 - (\lambda_1 + \lambda_2)\lambda + \lambda_1\lambda_2,$$

now by equality of polynomials we have that $a_1 = -(\lambda_1 + \lambda_2) < 0, \quad a_0 = \lambda_1\lambda_2 > 0$, and the inequalities *1* and *2* hold [3], for the inequality *3* we observe that if the polynomial of the second degree has two real, different roots then its determinant is positive $\Delta = a_1^2 - 4a_0 > 0$ i.e. the inequality *3* hold.

Conversely, if the condition (4) hold, the roots of polynomial $p(\lambda)$ are given by $\lambda = \dfrac{-a_1 \pm \sqrt{a_1^2 - 4a_0}}{2}$, the inequality 3 implies the existence of two different, real roots , the inequality 1 implies that $\dfrac{-a_1 + \sqrt{a_1^2 - 4a_0}}{2} > 0$, and the inequalities 1 and 2 implies that $\dfrac{-a_1 - \sqrt{a_1^2 - 4a_0}}{2} > 0$. ■

We are now ready to state and proof the main result of this section.

Theorem 1 *The monic polynomial of degree 3,* $p(\lambda) = \lambda^3 + a_2\lambda^2 + a_1\lambda + a_0$ *is a texture polynomial if and only if the following three conditions hold.*

1. $a_0, a_2 < 0 < a_1$.

2. $3a_1 < a_2^2$.

---

[3] In this work, a chain of inequalities $a < b < c < \ldots$, the first inequality is $a < b$, the second one is $b < c$ and so on.

3. If $\lambda_4 = \dfrac{-a_2 + \sqrt{a_2^2 - 3a_1}}{3}$ *and* $\lambda_5 = \dfrac{-a_2 - \sqrt{a_2^2 - 3a_1}}{3}$*, then* $p(\lambda_4) < 0$ *and* $p(\lambda_5) > 0$.

Remark. We observe that λ_4 and λ_5 are the roots of the first derivative of $p(\lambda)$, and therefore the condition 2 is equivalent to $\lambda_4, \lambda_5 \in \mathbb{R}$, and join to the condition 1 we have that $0 < \lambda_5 < \lambda_4$.

Proof. If $p(\lambda)$ is a texture polynomial, then exists three different $\lambda_i \in \mathbb{R}^+$, $(i = 1, 2, 3)$ such that $p(\lambda) = (\lambda - \lambda_1)(\lambda - \lambda_2)(\lambda - \lambda_3) = \lambda^3 - (\lambda_1 + \lambda_2 + \lambda_3)\lambda^2 + (\lambda_1\lambda_2 + \lambda_1\lambda_3 + \lambda_2\lambda_3)\lambda - \lambda_1\lambda_2\lambda_3$, by equality of polynomials we obtain

- $a_2 = -(\lambda_1 + \lambda_2 + \lambda_3) < 0,$
- $a_0 = -\lambda_1\lambda_2\lambda_3 < 0,$
- $a_1 = \lambda_1\lambda_2 + \lambda_1\lambda_3 + \lambda_2\lambda_3 > 0,$

and the condition *1* hold.

Without loss of generality, we suppose that $0 < \lambda_1 < \lambda_2 < \lambda_3$, if we have a polynomial of degree 3 with three real, simple roots then there exits two critical points, they are roots of the first derivative, i.e. the condition *2* hold. Now if the roots of polynomial are positive then the critical points are positive too and the follow chain of inequalities hold $\lambda_1 < \lambda_5 < \lambda_2 < \lambda_4 < \lambda_3$. From the coefficient of λ^3 is 1, we have that $\lim_{\lambda\to\infty} p(\lambda) = \infty$ and $\lim_{\lambda\to-\infty} p(\lambda) = -\infty$, then for points less than λ_1 the polynomial is negative, we applied the Rolle theorem to the roots λ_1 and λ_2, therefore the polynomial has a maximum value between λ_1 and λ_2, and therefore $p(\lambda_5) > 0$. Similarly for points greater than λ_3 the polynomial is positive and we applied the Rolle theorem to the roots λ_2 and λ_3, and the polynomial has a minimum, this value is negative i.e. $p(\lambda_4) < 0$.

Conversely, we have that $p(\lambda) = \lambda^3 + a_2\lambda^2 + a_1\lambda + a_0$ such that the conditions *1, 2* and *3* hold. The conditions *1,2* join to lemma 1 implies that $p'(\lambda)$ has two real, simple and positive roots given by :

$$\lambda_4 = \frac{-a_2 + \sqrt{a_2^2 - 3a_1}}{3}, \quad \lambda_5 = \frac{-a_2 - \sqrt{a_2^2 - 3a_1}}{3}.$$

First we observe that $0 < \lambda_5 < \lambda_4$, now computing $p''(\lambda_4) = 2\sqrt{a_2^2 - 3a_1} > 0$, this implies in λ_4 we have a minimum whereas $p''(\lambda_5) = -2\sqrt{a_2^2 - 3a_1} < 0$ and then in λ_5 we have a maximum. We applied repeatedly the Intermediate Value theorem. From *1*, we have that

$p(0) = a_0 < 0$ and from *3* it follows that $p(\lambda_5) > 0$, then we have a positive root. The condition *3*, $p(\lambda_4) < 0$ and $p(\lambda_5) > 0$, guarantee that exits a second root between λ_5 and λ_4, finally due to the coefficient to λ^3 is positive $p(\lambda)$ we have that $\lim_{\lambda\to\infty} = \infty$ and this implies $p(\lambda_4) < 0$, then $p(\lambda)$ intersects to horizontal axis one more time in the third root. ■

The condition *3*, of the theorem 1, can be replaced by

$$-2(a_2^2 - 3a_1)^{3/2} < 2a_2^3 - 9a_1a_2 + 27a_0 < 2(a_2^2 - 3a_1)^{3/2}, \tag{5}$$

the firs inequality is obtaining to solve $p(\lambda_5) > 0$ and the second one is obtained to solve $p(\lambda_4) < 0$. The condition (5) can be rewriting as

$$|2a_2^3 - 9a_1a_2 + 27a_0| < 2(a_2^2 - 3a_1)^{3/2}. \tag{6}$$

It is convenient to rewrite the theorem 1 in terms of the invariants of congruent matrices. This create directly a link between the matrix elements and its eigenvalues which facilitates subsequent computations and applications. To implement this fact, first we write the coefficients of its characteristic polynomial $p(\lambda)$ in terms of its trace ($\operatorname{tr} M$), trace of the square matrix ($\operatorname{tr} M^2$) and its determinant ($\det M$) in the following form:

$$p(\lambda) = \lambda^3 - \operatorname{tr} M\lambda^2 + \frac{1}{2}\left[\operatorname{tr}^2 M - \operatorname{tr} M^2\right]\lambda - \det M. \tag{7}$$

Now we are ready to present the main theorem of this section

Theorem 2 *A real, symmetric matrix M has its characteristic polynomial of textures, if and only if the following three conditions hold.*

1. *(a)* $\det M > 0$,

   (b) $\operatorname{tr} M > 0$,

   (c) $\operatorname{tr} M^2 < \operatorname{tr}^2 M$.

2. $\operatorname{tr}^2 M < 3\operatorname{tr} M^2$.

3. $|\operatorname{tr} M(5\operatorname{tr}^2 M - 9\operatorname{tr} M^2) - 54\det M| < \sqrt{2}(3\operatorname{tr} M^2 - \operatorname{tr}^2 M)^{3/2}$.

The theorems presented in this section will be applied to texture-zero matrices, the texture-zero formalism is the subject of next section.

IV. TEXTURE-ZERO FORMALISM

A texture-zero matrix is a 3×3 matrix with zeros in some entries, the way to count them is the following: a zero in the main diagonal add as 1, while zero off main diagonal add as 1/2. We need to sum all zeros for both mass matrices u-quarks and d-quarks. For example, given M_u and M_d as

$$M_u = \begin{pmatrix} * & 0 & * \\ 0 & * & * \\ * & * & 0 \end{pmatrix}, \qquad M_d = \begin{pmatrix} * & 0 & * \\ 0 & * & 0 \\ * & 0 & 0 \end{pmatrix}.$$

For M_u we have one zero in the main diagonal, we add (+1) and 2 zeros off main diagonal that add $1 = (1/2 + 1/2)$, then M_u has a 2-zero texture structure. Considering now M_d we have a 3-zero texture structure $(1 + 2)$. Then this set of matrices is said to have a 5-zero texture structure.

In this work we are going to consider a parallel structure for M_u and M_d mass matrices, this mean that if M_u has zeros in some places then M_d has zeros in the same position that M_u, because of that we will only have the following set of structures: 2-zero textures, 4-zero textures and 6-zero textures.

A. Notation

We start writing a symmetric matrix M given by:

$$M = \begin{pmatrix} E & D & F \\ D & C & B \\ F & B & A \end{pmatrix}.$$

This matrix is well determined by specifying six capital letters (A, B, C, D, E, F) and their corresponding positions, then we introduce the following notation:

- $M(x)$ is a matrix with a zero in the capital letter x, $(x = A, B, C, D, E, F)$.
- $M(x, y)$ is a matrix with zeros in the capital letters x and y, (x, y = A, B, C, D, E, F; $x \neq y$).
- $M(x, y, z)$ is a matrix with zeros in the capital letters x, y and z, $(x, y, z = A, B, C, D, E, F;\ x \neq y \neq z)$.

For example, a matrix with a zero in the position F is:

$$M(F) = \begin{pmatrix} E & D & 0 \\ D & C & B \\ 0 & B & A \end{pmatrix},$$

a matrix with zeros in the positions C and D is,

$$M(C, D) = \begin{pmatrix} E & 0 & F \\ 0 & 0 & B \\ F & B & A \end{pmatrix},$$

finally a matrix with zeros in the positions C, D and F is,

$$M(C, D, F) = \begin{pmatrix} E & 0 & 0 \\ 0 & 0 & B \\ 0 & B & A \end{pmatrix}.$$

Using this notation, we are able to list all possible textures, considering a parallel structure of mass matrices.

2-zero texture structure.

We have 6 different matrices, which are:

$M(A),\ M(C),\ M(E),\ M(B),\ M(D),\ M(F).$

4-zero texture structure.

In this case, we have 15 possibilities, which are:

$M(A, E),\ M(A, C),\ M(C, E),$

$M(A, B),\ M(A, D),\ M(A, F),$

$M(B, C),\ M(C, D),\ M(C, F),$

$M(B, E),\ M(D, E),\ M(E, F),$

$M(B, F),\ M(B, D),\ M(D, F).$

6-zero texture structure.

For this case, there are 20 different matrices, which are:

$M(A, B, C),\ M(A, C, F),\ M(A, C, D),$

$M(A, B, E),\ M(A, E, F),\ M(A, D, E),$

$M(B, C, E),\ M(C, E, F),\ M(C, D, E),$

$M(A, B, F),\ M(A, B, D),\ M(A, D, F),$

$M(B,C,F),\ M(B,C,D),\ M(C,D,F),$
$M(B,E,F),\ M(B,D,E),\ M(D,E,F),$
$M(A,C,E),\ M(B,D,F).$

Now we are ready to analyze which kind of textures have three different and positive eigenvalues, applying in each case one of the theorems presented in previous sections.

V. APPLICATIONS

In order to start the analysis in a systematic way, we need to implement another subclassification, which depends if the matrix has or not zeros in the main diagonal, doing this, first we analyze the 6-zeros textures, after this, we study the 4-zero textures and finally the 2-zeros textures.

A. 6-zeros analysis

According the sub-classification given above, the 6-zeros textures present the following cases:

1. Without zeros in the main diagonal there is one case $M(B,D,F)$.
2. With one zero in the main diagonal exist 9 cases: $M(A,B,F)$, $M(A,B,D)$, $M(A,D,F)$, $M(B,C,F)$, $M(B,C,D)$, $M(C,D,F)$, $M(B,E,F)$, $M(B,D,E)$, $M(D,E,F)$.
3. With two zeros in the main diagonal there are 9 cases: $M(A,B,C)$, $M(A,C,F)$, $M(A,C,D)$, $M(A,B,E)$, $M(A,E,F)$, $M(A,D,E)$, $M(B,C,E)$, $M(C,E,F)$, $M(C,D,E)$.
4. With three zeros in the main diagonal we have only 1 case ($M(A,C,E)$).

We obtain a total of 20 different possibilities. We only present the analysis of the following three cases.

- Applying the Theorem 2 (1b) the trivial $M(A,C,E)$ case is ruled out.
- Now, we analyze the Fritzsch 6-zero texture given by $M(C,E,F)$ [1]. Applying again the Theorem 2 (1b) we must have $\mathrm{tr}\, M(C,E,F) = A > 0$, from the condition (1a)

$\det M(C,E,F) = -D^2A < 0$ that is a contradiction, because of that this 6-zero texture is ruled out.

- Next, we analyze the following texture $M(A,D,F)$. The condition (1b) of the Theorem 2 we have that $\operatorname{tr} M(A,D,F) = C + E > 0$ and from (1a) $\det M(A,D,F) = -EB^2 > 0$, $\Leftrightarrow\ E < 0\ \Rightarrow\ C > 0\ \Rightarrow\ EC < 0$. The condition (1c) of the Theorem 2 implies that $0 < E^2 + C^2 + 2B^2 < E^2 + C^2 + 2EC\ \Rightarrow\ 0 < EC$ and we have a contradiction and this texture is also ruled out.

We have analyzed the others 17 cases and we found that the only case that is not excluded is $M(B,D,F)$, obviously being A, B and C the eigenvalues ($A \neq C \neq E > 0$).

B. 4-zeros analysis

These kind of textures have the following cases:

1. Without zeros in the main diagonal there are 3 cases: $M(B,F)$, $M(B,D)$, $M(D,F)$.
2. With one zero in the main diagonal exist 9 cases: $M(A,B)$, $M(A,D)$, $M(A,F)$, $M(B,C)$, $M(C,D)$, $M(C,F)$, $M(B,E)$, $M(D,E)$, $M(E,F)$
3. With two zeros in the main diagonal there are 3 cases: $M(A,E)$, $M(A,C)$, $M(C,E)$.

We present the analysis of some cases more representative:

- We start with the matrix $M(C,E)$. If we compute $\operatorname{tr}^2 M(C,E)$, $\operatorname{tr} M(C,E)^2$ and we apply the condition (1c) of the Theorem 2, we obtain:

  $$2(D^2 + F^2 + B^2) + A^2 < A^2,$$

  that is a contradiction. We have found that $M(A,E)$ and $M(A,C)$ are ruled out too.

- The second example is the Fritzsch 4-zero texture given by $M(E,F)$ [5]. From the Theorem 2 follows that the condition (1a) $\det M(E,F) = -AD^2 > 0$ implies $A < 0$, and of the condition (1b) $\operatorname{tr} M(E,F) = C + A > 0$ we have that $C > 0$ and then $AC < 0$. Now we compute $\operatorname{tr}^2 M(C,E)$, $\operatorname{tr} M(C,E)^2$ and using the condition (1c) of the Theorem 2, we obtain:

  $$2(D^2 + B^2) + C^2 + A^2 < C^2 + A^2 + 2AC,$$

then $AC > 0$, that is a contradiction.

We have analyzed the eight cases $M(A,B)$, $M(A,D)$, $M(A,F)$, $M(B,C)$, $M(C,D)$, $M(C,F)$, $M(B,E)$, $M(D,E)$ and we found that are ruled out.

The cases that are in agreement with the condition (1) of the Theorem 2 are $M(B,F)$, $M(B,D)$ and $M(D,F)$, this means that, these textures have possibilities to have real, positive and different eigenvalues. Now we find numerical values for the capital letters such that the conditions (2) and (3) hold. Now we present two numerical example for the u-quark mass matrix that satisfies Theorem 2. In the first example we have a small perturbation to the quark masses, and in the second one, we have different values.

$$M(B,F) = \begin{pmatrix} 3.00197 & 1.6 & 0 \\ 1.6 & 1300 & 0 \\ 0 & 0 & 174900 \end{pmatrix}, \qquad M(B,F) = \begin{pmatrix} 68.982 & 285 & 0 \\ 285 & 1234.02 & 0 \\ 0 & 0 & 174900 \end{pmatrix},$$

with eigenvalues[4] given by:

$$m_u = 3, \qquad m_c = 1300, \qquad m_t = 1749000.$$

in agreement with the experimental value [2].

C. 2-zeros analysis

Here we only have two cases,

1. Without zeros in the main diagonal belong three different possibilities $M(B)$, $M(D)$ and $M(F)$.

2. With one zero in the main diagonal also belong three different possibilities $M(A)$, $M(C)$ and $M(E)$.

We only present the analysis of $M(A)$. The condition (1b) produces $E + C > 0$, the condition (1a) implies that $2BDF - B^2E - F^2C > 0$ and the condition (1c) gives

---

[4] The units for all parameters are MeV, by they are omitted.

$2(B^2+D^2+F^2)+E^2+C^2 < E^2+C^2+2EC$, the last three inequalities are equivalents with

$$\begin{aligned} E+C &> 0, & (8)\\ 2BDF &> B^2E+F^2C, & (9)\\ 0 < B^2+D^2+F^2 &< EC, & (10) \end{aligned}$$

from (8) and (10) we have that $E>0$ and $C>0$, therefore

$$\begin{aligned} -2BF\sqrt{EC} &< B^2E+F^2C, & (11)\\ 2BF\sqrt{EC} &< B^2E+F^2C, & (12) \end{aligned}$$

now if $BF>0$, the inequalities (10, 12, 9) produce the following chain of inequalities

$$2BF\sqrt{B^2+D^2+F^2} < 2BF\sqrt{EC} < B^2E+F^2C < 2BDF,$$

and then

$$\sqrt{B^2+D^2+F^2} < D,$$

that is a contradiction. If $BF<0$ use (11). We have analyzed the other 2 cases $M(C)$, $M(E)$ and we found that are ruled out.

The cases that are in agreement with the condition (1) of the Theorem 2 are $M(B)$, $M(D)$ and $M(F)$. To finalize this section, we showing two numerical examples for 2-zero texture mass matrix for d-type quarks:

$$M(F)=\begin{pmatrix} 5.59533 & 3 & 0\\ 3 & 99.9056 & 2\\ 0 & 2 & 4210 \end{pmatrix}, \qquad M(F)=\begin{pmatrix} 9.94113 & 20 & 0\\ 20 & 95.9478 & 40\\ 0 & 40 & 4209.61 \end{pmatrix},$$

with eigenvalues given by:

$$m_u=5.5, \qquad m_c=100, \qquad m_t=4210.$$

VI. CONCLUSIONS

In this paper, in the understanding that by definition the physical mass of the quarks and charged leptons are the eigenvalues of the mass matrices. We found the necessary and sufficient conditions over the characteristic polynomial coefficients from any symmetric 3 by 3 matrix, so that it has real, simple and positive roots. We apply this formalism to analyze the symmetric texture-zero matrices. We found that a lot of them are ruled out when the first condition of the Theorem 2 is applied. Only the 4-zeros texture matrices $M(B,F)$, $M(B,D)$, $M(D,F)$ and the 2-zero texture matrices $M(B)$, $M(D)$, $M(F)$ are in agreement with this condition. In the texture-zero formalism, the matrices have variable coefficients, the conditions 2 and 3 impose restrictions over these coefficients, this means that, we need to find the complete domain of the coefficients in both mass matrices, u-type quarks and d-type quarks, in order to approximate the V_{ckm} matrix. Analytically this procedure is complicated, for this reason, we show numerical examples. Other restriction over the matrix coefficients is the V_{ckm} analysis of these kind of zero-textures, we shall discuss in detail this in a forthcoming paper [6].

Acknowledgments

A. Criollo acknowledges to Baltazar Aguirre Hernández for his useful comments. The authors also acknowledge to CONACyT(*México*), PROMEP(*México*) and SNI(*México*) for their financial support.

[1] H. Fritzsch, Phys. Lett. B **70**, 436 (1977). H. Fritzsch, Phys. Lett. B **73**, 317 (1978).

[2] J. Beringer *et al.* [Particle Data Group], Phys. Rev. D **86**, 010001 (2012). `http://pdg.lbl.gov/`

[3] H. Fritzsch and Z. z. Xing, Prog. Part. Nucl. Phys. **45**, 1 (2000) [arXiv:hep-ph/9912358].

[4] S. Friedberg, A. Insel and L. Spence. Linear Algebra. Fourth Edition. Prentice Hall (2006)

[5] H. Fritzsch and Z. -z. Xing, Phys. Lett. B **555**, 63 (2003) [hep-ph/0212195].

[6] A. Criollo *et al.* Work in progres.